

Initial Studies on the Contributions of Body Size and Gastrointestinal Passage Rates to Dietary Flexibility Among Gorillas

M.J. REMIS*

Department of Sociology and Anthropology, Purdue University,
West Lafayette, Indiana 47907-1365

KEY WORDS diet; digestive physiology; ecology; hominoid evolution; *Gorilla gorilla gorilla*

ABSTRACT Large body size has been traditionally seen as the primary dietary adaptation of gorillas, facilitating their consumption of fibrous foods (Schaller [1963] *The Mountain Gorilla*; Watts [1990] *Int. J. Primatol.* 11:21–45). Nevertheless, recent research has emphasized frugivory among western lowland gorillas, as well as the influence of habitat and seasonality on gorilla diet and behavior across subspecies (Watts [1990] *Int. J. Primatol.* 11:21–45; Tutin et al. [1991] *Philos. R. Soc. Trans. Lond. Biol.* 334:179–186; Remis [1994] Ph.D. Thesis, [1997a] *Am. J. Primatol.* 43:87–109, [1997b] *Am. J. Primatol.* 43:111–133, [1998] *Primate Locomotion: Recent Advances*, p 95–108, [1999] *Primates* 40:383–396; Nishihara [1995] *Primates* 36:151–168; Goldsmith [1999a] *Int. J. Primatol.* 20:1–23, [1999b] *Nonhuman Primates*, p 58–63). This study provides preliminary data to address the physiological underpinnings of dietary flexibility among gorillas, and their consumption of a broad range of fibrous and tannin-rich foods. To date, little is known about the digestive physiology of the African apes (but see Milton [1984] *Adaptations for Foraging in Nonhuman Primates*, p 249–279, Milton [1984][1999] *Evol. Anthropol.* 8:11–20; Milton and Demment [1988] *J. Nutr.* 118:1082–1088; Lambert [1997] Ph.D. Dissertation), although gastrointestinal morphology and proportions are roughly similar among species (Chivers and Hladik [1980] *J. Morphol.* 166:337–386). This study provides additional experimental data on the gastrointestinal passage times of gorillas (*Gorilla gorilla gorilla*) fed a captive diet in a zoological park setting and discusses results in relation to field research on gorilla feeding ecology. In this study, 480 small plastic markers were fed to six captive gorillas. The mean gut retention time (MRT) of the adult gorillas in this study was 50 hr, longer than the 31 hr reported for chimpanzees fed a similar diet (Lambert [1997] Ph.D. Dissertation). These data suggest that gorillas may retain foods in their gastrointestinal tracts longer than smaller hominoids, and that the large body size likely forms the primary basis for consumption of fiber. This research provides additional data to contribute to our understanding of the relationships of body size and morphology to ecology, and the evolution of body size, foraging strategy and social organization among the African apes. *Am J Phys Anthropol* 112:171–180, 2000. © 2000 Wiley-Liss, Inc.

Grant sponsor: National Science Foundation; Grant number: SBR-981584; Grant sponsor: Purdue University.

*Correspondence to: M.J. Remis, Department of Sociology and Anthropology, Purdue University, West Lafayette, IN 47907-1365.

E-mail: remis@sri.soc.purdue.edu

Received 20 July 1999; accepted 12 December 1999.

Gorillas have traditionally been classified as herbivore-folivores (Schaller, 1963; Fossey and Harcourt, 1977; Watts, 1990, 1996), and this diet has been seen as fundamental to cohesive grouping patterns (Wrangham, 1979). Nevertheless, gorillas' consumption of fruit and corresponding ranging, grouping, and social behavior vary with resource availability, within and between sites and subspecies (Tutin and Fernandez, 1985; Watts, 1990; Tutin et al., 1991, 1997; Remis, 1994, 1997a,b, 1998, 1999; Goldsmith, 1996, 1999a,b; Yamagiwa et al., 1996; Doran and McNeillage, 1998). At many sites, gorillas are seasonal frugivores, but they are often described as less persistent in their fruit-eating than sympatric chimpanzees (Nishihara, 1995; Kuroda et al., 1996; Yamagiwa et al., 1996) or bonobos (Malenky, 1990), and this factor likely shapes species-specific social and grouping behavior.

Generalizations concerning the relationship of body size to diet lead us to infer that gorillas should be more folivorous than sympatric chimpanzees (Gaulin and Konner, 1977; Temerin et al., 1984; Demment and Van Soest, 1985; Salier et al., 1985; Chivers and Langer, 1994). Moreover, the limited available physiological evidence suggests that gorillas may be anatomically equipped to digest fiber (Chivers and Hladik, 1980; Collet et al., 1984; Milton, 1984). As herbivores, high-altitude Virunga mountain gorillas conform to these expectations, but they live in an environment where fruits are scarce (Schaller, 1963; Watts, 1996). Virunga and Bwindi mountain gorillas have a more diverse diet and consume more fruit at lower elevations (McNeillage, 1995; Goldsmith, 1999b). Observational data on sympatric African apes in lowland forests reveal broad similarity and overlap in diets (Remis, 1997a; Kuroda et al., 1996; Tutin et al., 1991, 1997), which makes it difficult to use a size-based model to predict their diets (Schoener, 1971; Jarman, 1974; cf. Perrin, 1994).

Despite their penchant for fruit, the large size of gorillas probably lends them greater dietary flexibility than chimpanzees (Demment, 1983). It may facilitate niche separation during the dry season, when gorillas exploit fallback foods relatively high in

structural carbohydrates (fiber), condensed tannins, and phenols. On the other hand, during seasons of scarcity, chimpanzees rely on flexible grouping patterns to facilitate continued fruit consumption, although some of these fruits are fibrous or tannin-rich (Tutin and Fernandez, 1991, 1993; Wrangham and Waterman, 1983; Wrangham et al., 1991, 1993; Remis, 1997a, 1999; Remis, et al., unpublished findings; Reynolds et al., 1998).

DIETARY SELECTIVITY AND PLANT BIOCHEMISTRY

Variation in dietary selectivity among primates is presumed to reflect differences in digestive adaptations for metabolizing the nutrients and secondary compounds found in plants (Freeland and Janzen, 1974; Feeney, 1976; Rhodes and Cates, 1976; Glander, 1981; Waterman, 1984; Ganzhorn, 1989; Chivers and Langer, 1994; Milton, 1998). Folivorous primates often consume foods that are high in digestion inhibitors, such as lignin and secondary plant compounds (Oates et al., 1977; Freeland, 1991; Waterman and Kool, 1994), although many of these foods are also high in protein (Hladik, 1978; Milton, 1981; Mckey et al., 1981; Waterman et al., 1983; Davies and Oates, 1994).

While gorillas are commonly classified as folivores, they do not have the morphological specializations associated with forestomach fermentation that may allow colobines to detoxify alkaloids (Oates et al., 1977, 1980; Kay and Davies, 1994). Gorillas do, however, have a somewhat enlarged hindgut relative to chimpanzees and other frugivores, associated with high rates of colic-ecal fermentation (Eftman and Atkinson, 1950; Hosakawa and Kamiya, 1961; Bauchop, 1978; Parra, 1978; Chivers and Hladik, 1980; Clemmens and Philips, 1980; Milton and Demment, 1988). In addition, gorillas may use geophagia to facilitate consumption of secondary compounds (Watts, 1990; Remis, 1997a) and sometimes practice coprophagy under conditions of low food availability (Harcourt and Stewart, 1978; Hladik, 1978; Rogers et al., 1992).

Folivorous species generally have large populations of microbial gut flora and ap-

pear better able to digest high-fiber and tannin-rich foods than most frugivores (Milton, 1981; Milton and McBee, 1983; Watkins et al., 1985; Crissey et al., 1991; Dierenfeld et al., 1992). In fact, they might be committed to a digestive strategy (e.g., Nordin, 1981; Power and Oftedal, 1996) that requires them to consume leaves to meet their protein needs, even when fruit is abundant (Milton, 1981; Maisels, 1993; Tutin and Fernandez, 1994).

GORILLA BODY SIZE AND MORPHOLOGY: IMPLICATIONS FOR DIGESTIVE STRATEGIES AND DIET

Body size, in itself, has long been considered a mechanism for interspecific and intraspecific differences in diet (Bell, 1970; Jarman, 1974; Demment, 1983). Large body size facilitates digestion of high-fiber or chemically defended plant foods, and may allow gorillas to consume a folivorous diet without many of the gastrointestinal specializations of smaller-bodied folivores. The proportion of metabolic requirements to gut capacity decreases with body size. Larger animals can, therefore, retain foods longer in the gut and consume a lower-quality diet than smaller individuals with similar gut morphology (Demment and Van Soest, 1985; Feer, 1995; but see Lambert, 1997).

Gorillas are the largest and among the most sexually dimorphic of any primate species, with males typically twice the size of females (*G. g. gorilla* females approximately 72 kg, males 169 kg; Jungers and Susman, 1984; Zihlman, 1997). The evolution of differences in body size, muscle mass, and patterns of sexual dimorphism among the apes has been linked to species differences in socioecology, but these differences have yet to be fully evaluated (Leigh, 1994, 1995; Leigh and Shea, 1995; Janson and van Schaik, 1993; Martin et al., 1995; Zihlman and McFarland, unpublished findings).

In addition to body size, anatomical correlates of diet include craniodental and gastrointestinal specializations. All gorillas have a generalized craniodental morphology characteristic of the hominoids (Kay, 1975; Shea, 1983; Uchida, 1998; but see Hartman, 1988). Nevertheless, mountain gorillas have higher shearing crests and narrower central

incisors than western lowland gorillas, correlating with their greater folivory (Groves, 1986; Uchida, 1996). While gorillas are similar to chimpanzees in the morphology and general proportions of the gut (Chivers and Hladik, 1980, 1984; Martin et al., 1985), they have relatively larger colons with more cellulose-digesting ciliates (File et al., 1976; Collet et al., 1984). Preliminary investigations have suggested that gorillas have slow gut transit times relative to those of chimpanzees (Milton, 1984; but see Milton, 1999).

Retention of food in the gastrointestinal tract is presumed to allow animals to maximize the amount of nutrients absorbed (see Lambert, 1999 for review). Nevertheless, absolute surface area of the gut is likely as important as retention time in facilitating microbial activity and nutrient absorption (Kay and Davies, 1994). Large gut volume and surface area allow for greater production of volatile fatty acids as well as absorption and assimilation of these end products among simple stomached catarrhines (Bourton and Perrin, 1991). These features may permit gorillas to tolerate high levels of fiber, total phenols, and condensed tannins in their foods (Rogers et al., 1990; Cork and Foley, 1991). Gorillas may consume high-fiber foods in order to meet their total energetic or protein needs, to maximize digestive efficiency, or to maintain colonies of ciliates in the gut (Milton, 1984; Rogers et al., 1992; Tutin and Fernandez, 1994; Popovich et al., 1997).

The primary aim of this research was to conduct experimental observational study of gut passage times among captive gorillas. These initial laboratory-based data assist our efforts to better understand the effects of body size and digestive physiology on dietary flexibility of gorillas in their natural habitats.

METHODS

Study subjects, housing conditions, and diet

This study was conducted on six western lowland gorillas (*Gorilla gorilla gorilla*) housed at the San Francisco Zoo, March 1999. Study subjects included: three adult females, one of whom was lactating and one

TABLE 1. Study subjects Gorilla gorilla gorilla at the San Francisco Zoological Gardens

Individual	Sex	Age	Birth date	Weight in kg (date recorded)	History/condition
Pogo	F	41	1958	109.8 (3/96)	Wild-caught, hand-reared, nulliparous
Bawang	F	18	7/13/80	87 (4/96)	Cin Zoo born, hand-reared, lactating mother to female, born 9/98
Zura	F	17	9/13/81	77 (11/93)	Col. Zoo born, hand-reared, nulliparous
Mkubwa	M	23	5/01/75	169.6 (3/96)	San Francisco Zoo born, parent-raised
Shango	M	10	3/11/89	Est. 150	San Francisco Zoo born, mother-reared
Barney	M	5	10/01/93	Est. 50	San Francisco Zoo born, mother-reared

TABLE 2. Diet of study subjects at the San Francisco Zoological Gardens, 1999¹

Food	Male 4,900 kcal	Female 4,000 kcal
Banana	125 g	125 g
Apple	450 g	300 g
Orange	500 g	350 g
Yam	300 g	250 g
Carrot	900 g	675 g
Celery	3,000 g	1,500 g
Mixed greens (kale, collards, romaine)	500 g	500 g
Browse	200 g	200 g
High-Fiber Mazuri Leaf Eater Chow	800 g	400–700 g

¹In addition to the above set diets, large quantities of browse (*Acacia longifolia* branches), as well as six supplemental domestic vegetables per individual per day and scratch feed (2 parts cracked corn, 1 part popcorn, 1 part sunflower seeds, with additional grapes and peanuts) are scattered in the grotto. Animals consume these foods ad libitum. Milk or yogurt (8 oz) is also given daily to the lactating female. One and a half tablespoons of Metamucil mixed in water are also occasionally administered to Zura to facilitate the gut passage of foreign objects consumed in the grotto.

of whom was wild-born, a silverback, a blackback, and a juvenile male (see Table 1). The gorillas are currently group-housed in a large off-exhibit indoor night-quarter facility and have access to an outdoor grotto (22,000 square feet), weather permitting, from 10:00–16:00 hr daily. Mean temperature in the outdoor grotto was 20°C (range, 18.9–22.8°C), and the mean temperature indoors was 20.5°C (range, 16.7–25.0°C).

During AM and PM meals, at 8:00 and 16:00 hr, animals are temporarily separated into individual enclosures in the night quarters and fed meals of high-fiber monkey chow and fruits and vegetables (see Table 2). In addition, the animals receive browse, vegetables, grapes, and peanuts scattered in their outdoor grotto at 10 AM daily. This diet provides the gorillas approximately 17.85% dry-matter neutral detergent fiber and 9.46% dry-matter acid detergent fiber (Dier-

enfeld, personal communication). It is high-quality, relative to many of the foods consumed by gorillas in the wild (gorilla foods at Bai Hokou eaten during fruit scarcity: mean neutral detergent fiber = 61.6%, acid detergent fiber = 46.6%, Remis et al., unpublished findings).

No changes were made to the housing conditions or diet of the subjects during this study. Only the principal investigator and keepers were permitted in the area adjacent to the night quarters, as a health precaution, and to minimize any stress to the animals that might affect gut retention times.

Gut passage trials

Four experimental trials were conducted on each study animal. A trial consisted of the administration of 20 nontoxic colored plastic markers. The markers (measuring 4 × 2 × 1 mm) were concealed in a banana and offered to the subjects in an AM feed and PM feed. The methods, size, and type of marker were chosen to be directly comparable to the methods used during the study of gut passage rates of cercopithecines and chimpanzees by Lambert (1997). The markers are similar in size and weight to small seeds typically swallowed by monkeys and apes. Each animal was fed loaded bananas once in the morning and once in the late afternoon just before or after each feeding for a total of four doses of markers over 2 days. For all trials, each individual was assigned its own unique marker color.

Animals were observed from March 23, 1999 at 16:00 hr–March 29, 1999 at 8:00 AM, 136 hr after the initial AM trial. The gorillas were observed by two observers from 7:30–18:00 hr daily (with up to seven additional observers posted adjacent to the outdoor exhibit), and the time of all observed defeca-

tions was recorded. In addition, standardized estimates of defecation time were made by observing and gauging the dryness and hardness of several feces of known defecation time. Estimates of time of unobserved defecations were made using standardized means of fecal age based on appearance.

All fecal material was collected immediately or during enclosure cleaning, and washed and screened to determine whether colored markers were present. The number of markers and color(s) of markers were recorded. A total count of all defecations was maintained throughout the entire project. Three measures of gut passage were calculated. These include time of first marker appearance or transit time (TT), time of last appearance (TLA) of a marker, and mean retention time of markers (MRT). Mean retention time is considered to be the single best estimate of digesta movement through mammalian gastrointestinal tracts and is a measure of the average time of retention of all the elements of the digesta being studied. MRT is calculated:

$$t = \sum_{i=1} m_i t_i / \sum_{i=1} m$$

where m_i = the amount of marker excreted at the i th defecation at time t_i after dosing (Warner, 1981; Lambert, 1997).

RESULTS

Administration of markers

The marker-dosed bananas were generally readily and immediately consumed by the gorillas; no spitting of markers was observed. Animals did not typically detect the presence of the markers. In two cases, animals manually removed several markers from bananas; at least five markers fell on the cage floor during marker administration (6%) and were promptly removed from the enclosure. Thus, the gorillas apparently swallowed a total of 475 markers during this study.

Defecation of markers

The gorillas defecated 193 times between March 24, 1999 at 8:00 hr and March 28, 1999 at 16:00 hr, resulting in an average of 7.3 defecations per individual per 24-hr day.




Fig. 1. Passage of markers fed to gorillas at the San Francisco Zoo, 1999.

On average, 1.6 markers were excreted per individual defecation (range, 0–8). The number of different color markers per defecation ranged from 1–4, so considerable mixing of the gut contents occurred. A total of 309 markers was excreted and collected (65% of 475 swallowed) (Fig. 1). As in other studies, a considerable number of markers were not recovered during the study period (Warner, 1981; Lambert, 1997). Some unrecovered markers may have been overlooked in the fibrous dung during the screening process. Other markers may have temporarily adhered to the intestinal walls; a small number of markers were still appearing in stools, albeit infrequently, at the end of the study period. Gorilla feces were typically large, fibrous, and well-formed. However, on the evening following administration of the first two trials of markers, the silverback produced a single defecation of loose stool when he responded aggressively with chases and bark vocalizations to a brief challenge from the large blackback for first access to the door to the indoor quarters. The conflict may have influenced the relatively rapid transit time of both males (see below). Thus, gut transit and retention times reported here should likely be considered as minimum estimates.

TABLE 3. *Gastrointestinal passage times in hours of gorillas at the San Francisco Zoological Gardens*

Individual	Age/sex	Body weight (kg)	Percentage of markers recovered	Transit time	Mean rate of transit	Time of last appearance of marker
Pogo	Ad F	109	59	36.0	56.6	96.0
Bawang	Ad F	87	53	24.0	54.7	120.0
Zura	Ad F	77	41	31.2	61.9	136.0
Mkubwa	Ad M	170	54	17.0	38.6	112.0
Shango	SAd M	150	100	22.0	43.4	96.0
Barney	J M	50	79	8.0	36.3	70.0
Group Mean		110	64	23.0	47.1	109.3
Adult Mean		118	61	25.9	50.0	112.0

Gut passage time

The gorillas in this study exhibited considerable individual variation (see also Warner, 1981) in transit time (TT), mean gut passage time, i.e., mean rate of transit (MRT), and time of last appearance of markers (TLA) (ANOVA MRT, $F = 15.52$, $P = 0.000$) (Table 3). MRT of markers also varied by trial (ANOVA $F = 4.162$, $P < 0.007$), and may have been affected by minor dietary variation. MRT for all trials for the adult gorillas in this study was 50 hr (range, 16.5–136 hr). As expected, the juvenile gorilla had the shortest MRT (36.3 hr, $n = 63$ markers). The silverback and blackback did not differ in MRT (independent samples t -test $t = -1.5$, $P < 0.133$). As a result, data for the silverback and blackback were combined and data on adult-sized animals were analyzed for sex differences. Contrary to body-size expectations, the MRT for the large males (MRT = 41.74, $n = 123$ markers) was shorter than the MRT for the three females in this study (MRT = 57.38, $n = 122$ markers, independent samples t -test, $t = -5.3$, $P = 0.000$).

DISCUSSION

The five adult-sized gorillas in this study had similar mean transit times and mean retention times to those recently reported for other captive gorillas, chimpanzees, and orangutans (Caton, 1999; Milton, 1999), suggesting that gut retention may be a phylogenetically determined trait. Nevertheless, the San Francisco Zoo gorillas had longer mean gut retention times and mean time of last appearance of markers than the two adult chimpanzees studied by Lambert (TT = 23.2, MRT = 31.5, TLA = 63.3 hr). This difference may be significant, as this

study was more comparable in methods and diet to the study by Lambert, 1997 (chimpanzee diet consisted of 700 g Mazuri Leaf Eater Chow and 1.8 kg mixed produce daily) than to those described in other studies. The single juvenile gorilla in this study was similar in both body weight (50 kg) and mean retention time (MRT = 36 hr) to the adult chimpanzees in Lambert's study.

Dietary intake, the proportion of fiber in the diet of captive animals, feeding frequency, ambient temperature, pregnancy, activity level, age, and exposure to medications which could destroy natural gut fauna and flora all affect gut passage rates (Warner, 1981; Milton, 1999). Our efforts to understand the digestive physiology of hominoids are just beginning, and sample sizes remain small. Therefore, the relative contributions of ecological niche, phylogeny, body size, sex, and dietary intake to gut retention have yet to be determined (Lambert, 1997; Milton, 1999). Future research will seek to expand the sample size of this study to include chimpanzees and additional gorilla subjects. It will also investigate the effects of an ad libitum all-fruit (low-fiber) or high-fiber diet on gut retention times. This research should provide clues as to whether seasonal dietary flexibility among gorillas corresponds to changes in digestive strategies and efficiency, as has been found for other taxa (Gross et al., 1986; Lee and Houston, 1993; Afik and Karasov, 1995; Afik et al., 1997; cf. Milton, 1999).

The results of this study confirm expectations of the effects of body size on ape gut retention times, but not expectations of body-size-related sex differences in gorilla MRT. The male and female subjects in this study were fed different amounts of fruit

and chow, although vegetables and browse were consumed *ad libitum*. Differences in fixed-intake as well as the social factors discussed above may have affected sex differences in MRT.

Between species, the large size of gorillas permits a high-fiber diet, relative to smaller-bodied chimpanzees, although more work on both species at various intake rates is warranted. Gorillas may also retain foods longer than chimpanzees fed similar diets. Nevertheless, relative to body size, great apes have shorter gut passage times than might be expected based on the gut passage times of much smaller-bodied cercopithecines and the morphologically specialized colobine folivores (Lambert, 1997; Milton, 1999). These data support the hypothesized divergence between ape and monkey digestive strategies and corresponding differences in dietary quality relative to body weight (Andrews, 1981; Temerin and Cant, 1983; Lambert, 1997; Wrangham et al., 1998). Proportionately longer transit times appear to represent an alternative means by which small simple-stomached catarrhines consume foods containing high levels of fiber and secondary compounds (Lambert, 1997, 1999).

Gorillas eat large quantities of fleshy fruit across study sites, but rely on fibrous fruits and foliage as staples or fallbacks (Williamson et al., 1990). The nutritional profiles of gorillas reveal selection of foods high in proteins or sugar relative to fiber and tannins (Casimir, 1975; Goodall, 1977; Calvert, 1985; Rogers et al., 1990; Watts, 1990; Plumptre, 1995; Remis et al., unpublished findings). Consumption of tannins varies with habitat and presumed variability of investment of plants in secondary compounds (see also Reynolds et al., 1998).

Wrangham et al. (1998) demonstrated that, despite body size differences, the nutrient and biochemical profiles of foods eaten by sympatric primates at Kibale were similar (see also Conklin et al., 1998). Likewise, chemical analyses of fruits consumed by gorillas, chimpanzees, and grey-cheeked mangabeys at Bai Hokou, Central African Republic (CAR) during a poor fruit season show similar site-specific biochemical profiles of fruits consumed by all species (Remis et al., unpublished findings). Large body

size and surface area of the gut and long retention of foods in the gastrointestinal tract may work together, or separately, to facilitate dietary flexibility and the consumption of foods containing digestion inhibitors.

This study provides further preliminary insights into the physiological bases of food choice among gorillas. Gorillas share a generalized hominoid morphology and physiology and a penchant for fleshy fruit with the smaller-bodied apes. Nevertheless, gorillas' body size lends them flexibility to subsist on alternate foods as staples or fallbacks whenever necessary. Large body size also spells some disadvantages for gorillas, as they need absolutely large amounts of foods and occasionally face constraints on their access to arboreal resources (Remis, 1995, 1999).

Research on dietary strategies and physiology also provides insight into the possible ecological factors mediating species diversification among the African apes and between the African apes and monkeys. For example, during the evolutionary diversification of the African hominoids, increasing body size from a smaller highly frugivorous ancestor probably lent gorillas dietary flexibility. Consumption of fibrous fallback or staple foods likely facilitated gorilla group cohesion (Wrangham, 1979; Remis, 1997a,b) as well as coexistence with sympatric chimpanzees, and permitted gorillas to expand their range or retreat into high-altitude areas where fruit is scarce (Groves, 1986; Yamagiwa et al., 1996).

ACKNOWLEDGMENTS

This project received the approval of the Purdue Animal Use and Care Committee. The curators, veterinarians, and staff at the San Francisco Zoo, especially E. Sargent, E. Lyon, and F. Dunker, facilitated this project. This study would not have been possible without the avid participation of keeper Mary Kerr. Kerr and other keepers, San Francisco Zoo docent volunteers, and students from the University of California at Santa Cruz graciously assisted in the recording of behavior and labor-intensive searching for markers in fecal material. Additional thanks go to the curious and cooperative study subjects who assisted in the

production and collection of samples. E. Dierenfeld and J. Lambert provided assistance on the design of this project. The comments of R. McFarland, M. Kerr, and three reviewers improved the manuscript.

LITERATURE CITED

- Afik D, Karasov WH. 1995. The tradeoffs between digestion rate and efficiency in warblers and their ecological implications. *Ecology* 76:2247–2257.
- Afik D, Darken BW, Karasov WH. 1997. Is diet shifting facilitated by modulation of intestinal nutrient uptake? A test of an adaptational hypothesis in yellow-rumped warblers. *Phys Zool* 70:213–221.
- Andrews P. 1981. Species diversity and diet in monkeys and apes during the Miocene. In: Stringer CB, editor. *Aspects of human evolution*. London: Taylor and Francis. p 25–61.
- Bauchop T. 1978. Digestion of leaves in vertebrate arboreal folivores. In: Montgomery GG, editor. *The ecology of arboreal folivores*. Washington, DC: Smithsonian Institution Press. p 193–204.
- Bell RH. 1970. A grazing ecosystem in the Serengeti. *Sci Am* 225:155–170.
- Bourton MR, Perrin MR. 1991. Comparative gut morphometrics of Vervet (*Cercopithecus aethiops*) and Samango (*C. mitis erytharchus*) monkeys. *Z Saugtierkunde* 56:65–71.
- Calvert JJ. 1985. Food selection by western gorillas (*Gorilla gorilla gorilla*) in relation to food chemistry. *Oecologia* 65:236–246.
- Casimir MJ. 1975. Feeding ecology and nutrition of an eastern gorilla group in the Mt. Kahuzi region (Republic of Zaire). *Folia Primatol (Basel)* 24:81–136.
- Caton JM. 1999. A preliminary report on the digestive strategy of the western lowland gorilla. *Aust Primatol* 13:2–7.
- Chivers DJ, Hladik CM. 1980. Morphology of the gastrointestinal tract in primates: comparisons with other mammals in relation to diet. *J Morphol* 166:337–386.
- Chivers DJ, Hladik CM. 1984. Diet and gut morphology in primates. In: Chivers DJ, Wood B, Bilsborough A, editors. *Food acquisition and processing in primates*. New York: Plenum Press. p 213–230.
- Chivers DJ, Langer P, editors. 1994. *The digestive system in mammals: food form and function*. Cambridge: Cambridge University Press. 446 p.
- Clemmens ET, Phillips B. 1980. Organic acid production and digesta movement in the gastrointestinal tract of the baboon and sykes monkey. *Comp Biochem Physiol* 66:529–532.
- Collet JY, Bourreau E, Cooper RW, Tutin CEG, Fernandez M. 1984. Experimental demonstration of cellulose digestion by *Troglodytes gorillae*, an intestinal ciliate of lowland gorillas. *Int J Primatol* 5:328.
- Conklin NL, Wrangham RW, Hunt KD. 1998. Dietary response of chimpanzees and cercopithecines to seasonal variation in fruit abundance. II. Macronutrients. *Int J Primatol* 19:971–998.
- Cork SJ, Foley WJ. 1991. Digestive and metabolic strategies of arboreal folivores in relation to chemical defenses in temperate and tropical forests. In: Palo RT, Robbins CT, editors. *Plant defenses against mammalian herbivory*. Boca Raton: CRC Press. p 133–188.
- Crissey SD, Edwards MS, Oftedal OT, Currier JA, Rudran R. 1991. The role of fiber in natural and manufactured diets fed to red howler monkeys (*Alouatta seniculus*). In: Meehan TP, Thompson SD, Allen ME, editors. *Proceedings of the Eighth Dr. Scholl Conference on the Nutrition of Captive Wild Animals*. Chicago: Lincoln Park Zoological Society. p 213–230.
- Davies AG, Oates JF. 1994. *Colobine monkeys: their ecology, behavior and evolution*. Cambridge: Cambridge University Press. 415 p.
- Demment MW. 1983. Feeding ecology and the evolution of body size in baboons. *Afr J Ecol* 21:219–233.
- Demment MW, van Soest PJ. 1985. A nutritional explanation of body-size patterns of ruminant and non-ruminant herbivores. *Am Nat* 125:641–672.
- Dierenfeld ES, Koontz FW, Goldstein RS. 1992. Feed intake, digestion and passage of the Proboscis monkey (*Nasalis larvatus*) in captivity. *Primates* 33:399–405.
- Doran DM, McNeillage A. 1998. Variation in behavior of gorilla subspecies: what's diet got to do with it? *Evol Anthropol* 6:120–131.
- Eftman H, Atkinson WB. 1950. Abdominal viscera of the gorilla. In: Gregory WK, editor. *The anatomy of the gorilla*. New York: Columbia University Press. p 197–205.
- Feeney PP. 1976. Plant apparency and chemical defence. *Rec Adv Phytochem* 10:1–40.
- Feer F. 1995. Seed dispersal in African forest ruminants. *J Trop Ecol* 11:683–689.
- File SK, McGrew WC, Tutin CEG. 1976. The intestinal parasites of a community of feral chimpanzees, *Pan troglodytes schweinfurthii*. *J Parasitol* 62:259–161.
- Fossey D, Harcourt AH. 1977. Feeding ecology of free-ranging mountain gorilla (*Gorilla gorilla beringei*). In: Clutton-Brock TH, editor. *Primate ecology*. London: Academic Press. p 441–464.
- Freeland WJ. 1991. Plant secondary metabolites: biochemical coevolution with herbivores. In: Palo RT, Robbins CT, editors. *Plant defenses against mammalian herbivory*. Boston: CRC Press. p 61–81.
- Freeland WJ, Janzen DH. 1974. Strategies in herbivory by mammals: the role of plant secondary compounds. *Am Nat* 108:269–289.
- Ganzhorn JU. 1989. Primate species separation in relation to secondary plant chemicals. *Hum Evol* 4:125–132.
- Gaulin SJC, Konner M. 1977. On the natural diet of primates including humans. In: Wurtman RJ, Wurtman JJ, editors. *Nutrition and the brain*. New York: Raven Press. p 1–86.
- Glander KE. 1981. The impact of plant secondary compounds on primate feeding behavior. *Yrbk Phys Anthropol* 25:1–18.
- Goldsmith ML. 1996. Ecological influences on the ranging and grouping behavior of western lowland gorillas at Bai Hokou, Central African Republic. Ph.D. dissertation. Stony Brook: State University of New York at Stony Brook.
- Goldsmith ML. 1999a. Ecological constraints on the foraging effort of western gorillas (*Gorilla gorilla gorilla*) at Bai Hokou, Central African Republic. *Int J Primatol* 20:1–23.
- Goldsmith ML. 1999b. Gorilla socioecology. In: Dolhinov P, Fuentes A, editors. *The nonhuman primates*. Mountain View, CA: Mayfield Publishing. p 58–63.
- Goodall AG. 1977. Feeding and ranging of a mountain gorilla (*Gorilla g. beringei*) group in the Tshibinda-Kahuzi region (Zaire). In: Clutton-Brock TW, editor. *Primate ecology*. New York: Academic Press. p 450–479.
- Gross J, Wang Z, Wunder BA. 1986. Adaptations to food quality and energy needs: changes in gut morphology and capacity of *Microtus orchrogaster*. *J Mammal* 66:661–669.
- Groves CP. 1986. Systematics of the great apes. In: Swindler DR, Erwin J, editors. *Comparative primate*

- biology, volume 1. Systematics, evolution and anatomy. New York: Alan R. Liss. p 187–217.
- Harcourt AH, Stewart KJ. 1978. Coprophagy by wild mountain gorilla. *E Afr Wild J* 16:223–225.
- Hartman JE. 1988. A cladistic analysis of hominoid molars. *J Hum Evol* 17:489–502.
- Hladik CM. 1978. Adaptive strategies of primates in relation to leaf-eating. In: Montgomery GG, editor. *The ecology of arboreal folivores*. Washington, DC: Smithsonian Institution Press. p 373–395.
- Hosakawa H, Kamiya T. 1961. Anatomical sketches of the visceral organs of the mountain gorilla (*Gorilla gorilla beringei*). *Primates* 3:1–21.
- Janson CH, van Schaik CP. 1993. Ecological risk aversion in juvenile primates: slow and steady wins the race. In: Pereira ME, Fairbanks LA, editors. *Juvenile primates*. New York: Oxford University Press. p 57–76.
- Jarman PJ. 1974. The social organization of antelope in relation to their ecology. *Behavior* 48:215–267.
- Jungers WL, Susman RL. 1984. Body size and skeletal allometry in African apes. In: Susman RL, editor. *The pygmy chimpanzee: evolutionary biology and behavior*. New York: Plenum Press. p 233–273.
- Kay RNB. 1975. The functional adaptations of primate molar teeth. *Am J Phys Anthropol* 43:195–216.
- Kay RNB, Davies AG. 1994. Digestive physiology. In: Davies AG, Oates JF, editors. *Colobine monkeys, their ecology behaviour and evolution*. Cambridge: Cambridge University Press. p 229–250.
- Kuroda S, Nishihara T, Suzuki S, Oko RA. 1996. Sympatric chimpanzees and gorillas in the Ndoki Forest, Congo. In: McGrew WC, Marchant L, Nishida T, editors. *Great ape societies*. Cambridge: Cambridge University Press. p 71–81.
- Lambert JE. 1997. Digestive strategies, fruit processing, and seed dispersal in the chimpanzees (*Pan troglodytes*) and redtail monkeys (*Cercopithecus ascanius*) of Kibale National Park, Uganda. Ph.D. dissertation. Urbana, IL: University of Illinois at Urbana-Champaign.
- Lambert J. 1999. Primate digestion: interactions among anatomy physiology, and feeding ecology. *Evol Anthropol* 7:8–20.
- Lee WB, Houston DC. 1993. The effects of diet quality on gut anatomy in British voles. *J Comp Physiol [B]* 163:337–339.
- Leigh SR. 1994. Ontogenetic correlates of diet in anthropoid primates. *Am J Phys Anthropol* 94:499–522.
- Leigh SR. 1995. Socioecology and the ontogeny of sexual size dimorphism in anthropoid primates. *Am J Phys Anthropol* 97:339–356.
- Leigh SR, Shea BT. 1995. Ontogeny and the evolution of adult body size dimorphism in apes. *Am J Primatol* 36:37–60.
- Maisels F. 1993. Gut passage rate in guenons and mangabeys: another indicator of a flexible dietary niche? *Folia Primatol (Basel)* 61:35–37.
- Malenky RK. 1990. Ecological factors affecting food choices and social organization in *Pan paniscus*. Ph.D. dissertation. Stonybrook, NY: State University of New York at Stonybrook.
- Martin RD, Chivers DJ, Maclarnon AM, Hladik CM. 1985. Gastrointestinal allometry in primates and other mammals. In: Jungers WL, editor. *Size and scaling in nonhuman primates*. New York: Plenum Press. p 61–89.
- Martin RD, Wilner LA, Dettling A. 1995. The evolution of sexual size differences in primates. In: Short RV, Balaban E, editors. *The differences between the sexes*. Cambridge: Cambridge University Press. p 159–200.
- McKey DB, Gartlan JS, Waterman PG, Choo GH. 1981. Food selection by black colobus monkeys (*Colobus satanas*) in relation to plant chemistry. *Biol J Linn Soc* 16:115–146.
- McNeillage AJ. 1995. Mountain gorillas in the Virunga volcanoes: ecology and carrying capacity. Ph.D. dissertation. Bristol, UK: University of Bristol.
- Milton K. 1981. Food choice and digestive strategies of two sympatric primate species. *Am Nat* 117:496–505.
- Milton K. 1984. The role of food-processing factors in primate food choice. In: Rodman PS, Cant JGH, editors. *Adaptations for foraging in nonhuman primates*. New York: Columbia University Press. pp 249–279.
- Milton K. 1998. Physiological ecology of howlers (*Alouatta*): energetic and digestive considerations and comparison with the colobinae. *Int J Primatol* 19: 513–548.
- Milton K. 1999. A hypothesis to explain the role of meat-eating in human evolution. *Evol Anthropol* 8:11–20.
- Milton K, Demment MW. 1988. Digestion and passage kinetics of chimpanzees fed high and low fiber diets and comparison with human data. *J Nutr* 118:1082–1088.
- Milton K, McBee RH. 1983. Structural carbohydrate digestion in a New World primate, *Alouatta palliata* Gray. *Comp Biochem* 74:29–31.
- Mole S, Waterman PG. 1987. Tannins as antifeedants to mammalian herbivores, still an open question? In: Waller GR, editor. *Allochemicals: role in agriculture and forestry*. Washington, DC: American Chemical Society. p 572–587.
- Nishihara T. 1995. Feeding ecology of western lowland gorillas in the Nouabale-Ndoki National Park, Congo. *Primates* 36:151–168.
- Nordin M. 1981. Voluntary food intake and digestion in Malaysian primates with special reference to the intake of energy and protein. *Malays Appl Biol* 10:163–175.
- Oates JF, Swain T, Zantovska J. 1977. Secondary compounds and food selection by colobus monkeys. *Biochem Syst Ecol* 5:317–321.
- Oates JF, Waterman PG, Choo GM. 1980. Food selection by the south Indian leaf-monkey, *Presbytis johinii*, in relation to leaf chemistry. *Oecologia* 45:45–56.
- Parra R. 1978. Comparison of foregut and hindgut fermentation in herbivores. In: Montgomery GG, editor. *The ecology of arboreal folivores*. Washington, DC: Smithsonian Press. p 205–229.
- Perrin MR. 1994. Herbivory and niche partitioning. In: Chivers DJ, Langer P, editors. *The digestive system in mammals: food, form and functioning*. Cambridge: Cambridge University Press. p 128–149.
- Plumptre AJ. 1995. The chemical composition of montane plants and its influence on the diet of large mammalian herbivores in the Parc National des Volcans, Rwanda. *J Zool Lond* 235:323–337.
- Popovich DG, Jenkins DJA, Kendall CWC, Dierenfeld ES, Carroll RW, Tariq N, Vidgen E. 1997. The western lowland gorilla diet has implications for the health of humans and other hominoids. *J Nutr* 127: 2000–2006.
- Power ML, Oftedal OT. 1996. Differences among captive callitrichids in the digestive responses to dietary gum. *Am J Primatol* 40:131–144.
- Remis MJ. 1994. Feeding ecology and positional behavior of western lowland gorillas (*Gorilla gorilla gorilla*) in the Central African Republic. Ph.D. thesis. New Haven: Yale University.
- Remis MJ. 1995. The effects of body size and social context on the arboreal activities of lowland gorillas (*Gorilla gorilla gorilla*) in the Central African Republic. *Am J Phys Anthropol* 97:413–433.
- Remis MJ. 1997a. Gorillas as seasonal frugivores: use of resources that vary. *Am J Primatol* 43:87–109.
- Remis MJ. 1997b. Ranging and grouping patterns of a western lowland gorilla (*Gorilla gorilla gorilla*) group

- at Bai Hokou, Central African Republic. *Am J Primatol* 43:111–133.
- Remis MJ. 1998. The gorilla paradox: effects of body size and habitat on positional behavior of lowland and mountain gorillas. In: Strasser E, Fleagle JG, Rosenberger AL, McHenry HM, editors. *Primate locomotion: recent advances*. New York: Plenum Press. p 95–108.
- Remis MJ. 1999. Tree structure and sex differences in arboreality among lowland gorillas at Bai Hokou, Central African Republic. *Primates* 40:383–396.
- Remis M, Dierenfeld ES, Mowry CB, Carroll RW. Unpublished findings. Nutritional aspects of western lowland gorilla (*Gorilla gorilla gorilla*) diet during seasons of fruit scarcity at Bai Hokou, Central African Republic. *Oecologia*.
- Reynolds V, Plumptre AJ, Greenham J, et al. 1998. Condensed tannins and sugars in the diet of chimpanzees (*Pan troglodytes schweinfurthii*) in the Budongo Forest, Uganda. *Oecologia* 115:331–336.
- Rhodes DF, Cates RG. 1976. Toward a general theory of plant anti-herbivore chemistry. In: Wallace JW, Mansell RL, editors. *Biochemical interaction between plants and insects*. New York: Plenum Press. p 168–213.
- Rogers ME, Williamson EA, Tutin CEG, Fernandez M. 1990. Gorilla diet in the Lope Reserve, Gabon: a nutritional analysis. *Oecologia* 84:326–339.
- Rogers ME, Maisels F, Williamson EA, Tutin CEG, Fernandez M. 1992. Nutritional aspects of gorilla food choice in the Lope Reserve, Gabon. In: Itoigawa N, Sugiyama Y, Sackett GP, Thompson RKR, editors. *Topics in primatology, volume 2: behavior, ecology and conservation*. Tokyo: University of Tokyo Press. p 385–394.
- Salier LD, Gaulin JC, Boster JS, Kurland JA. 1985. Measuring the relationship between dietary quality and body size in primates. *Primates* 26:14–27.
- Schaller GB. 1963. *The mountain gorilla: ecology and behavior*. Chicago: University of Chicago Press.
- Schoener TW. 1971. Theories of feeding strategies. *Annu Rev Ecol System* 2:369–404.
- Shea BT. 1983. Size and diet in the evolution of African ape craniodental form. *Folia Primatol (Basel)* 40:32–68.
- Temerin AB, Wheatley BS, Rodman PS. 1984. Body size and foraging in primates. In: Rodman PS, Cant JGH, editors. *Adaptations for foraging in nonhuman primates. Contributions to an organismal biology of prosimians, monkeys and apes*. New York: Columbia University Press. p 217–248.
- Temerin AL, Cant JGH. 1983. The evolutionary divergence of Old World monkeys and apes. *Am Nat* 122: 335–351.
- Tutin CEG, Fernandez M. 1985. Foods consumed by sympatric populations of *Gorilla g. gorilla* and *Pan t. troglodytes* in Gabon: some preliminary data. *Int J Primatol* 6:27–43.
- Tutin CEG, Fernandez M. 1991. Responses of wild chimpanzees and gorillas to the arrival of primatologists: behavior observed during habituation. In: Box HO, editor. *Primate responses to environmental change*. London: Chapman and Hall. p 187–197.
- Tutin CEG, Fernandez M. 1993. Composition of the diet of chimpanzees and comparisons with that of sympatric lowland gorillas in the Lope Reserve, Gabon. *Am J Primatol* 30:195–211.
- Tutin CEG, Fernandez M. 1994. Comparison of food processing by sympatric apes in the Lope Reserve, Gabon. In: Thierry B et al., editors. *Current primatology volume 1: ecology and evolution*. Strasbourg, France: University Louis Pasteur.
- Tutin CEG, Fernandez M, Rogers ME, Williamson EA, McGrew WC. 1991. Foraging profiles of sympatric lowland gorillas and chimpanzees in the Lope Reserve, Gabon. *Philos Trans R Soc Lond Biol* 334:179–186.
- Tutin CEG, Ham RM, White LJT, Harrison MJS. 1997. The primate community of the Lope Reserve, Gabon: diets, responses to fruit scarcity and effects on biomass. *Am J Primatol* 42:1–24.
- Uchida A. 1996. What we don't know about great ape variation. *Trends Evol Ecol* 11:163–168.
- Uchida A. 1998. Variation in gorilla dental morphology. *J Hum Evol* 34:55–70.
- Warner AC. 1981. Rate of passage of digesta through the gut of mammals and birds. *Nutr Abstr Rev Ser B* 51:789–820.
- Waterman PG. 1984. Food acquisition and processing as a function of plant chemistry. In: Chivers DJ, Wood B, Bilsborough A, editors. *Food acquisition and processing in primates*. New York: Plenum Press. p 177–211.
- Waterman PG, Kool K. 1994. Colobine food selection and plant chemistry. In: Davies AG, Oates JF, editors. *Colobine monkeys: their ecology, behavior and evolution*. Cambridge: Cambridge University Press. p 251–284.
- Waterman PG, Choo GM, Vedder AL, Watts DP. 1983. Digestibility, digestion-inhibitors, and nutrients of herbaceous foliage and green stems from an African montane flora and comparison with other tropical flora. *Oecologia* 60:244–249.
- Watkins BE, Ullrey DE, Whetter PA. 1985. Digestibility of a high-fiber biscuit-based diet by black and white colobus (*Colobus guereza*). *Am J Primatol* 9:137–144.
- Watts DP. 1990. Ecology of gorillas and its relation to female transfer in mountain gorillas. *Int J Primatol* 11:21–45.
- Watts DP. 1996. Comparative socio-ecology of gorillas. In: McGrew WC, Marchant LF, Nishida T, editors. *Great ape societies*. Cambridge: Cambridge University Press. p 16–28.
- Williamson EA, Tutin CEG, Rogers ME, Fernandez M. 1990. Composition of the diet of lowland gorillas at Lope Gabon. *Am J Primatol* 21:265–277.
- Wrangham RW. 1979. On the evolution of ape social systems. *Soc Sci Inf* 18:334–368.
- Wrangham RW, Waterman PG. 1983. Condensed tannins in fruits eaten by chimpanzees. *Biotropica* 15: 217–222.
- Wrangham RW, Conklin NL, Chapman CA, Hunt KD. 1991. The significance of fibrous foods for Kibale Forest chimpanzees. *Philos Trans R Soc Lond Biol* 334: 11–18.
- Wrangham RW, Conklin NL, Etot G, Obua J, Hunt KD, Hauser MD, Clark AP. 1993. The value of figs to chimpanzees. *Int J Primatol* 14:243–256.
- Wrangham RW, Conklin NL, Hunt KD. 1998. Dietary response of chimpanzees and cercopithecines to seasonal variation in fruit abundance. I. Antifeedants. *Int J Primatol* 19:949–970.
- Yamagiwa J, Maruhashi T, Yumoto T, Mwanza N. 1996. Dietary and ranging overlap in sympatric gorillas and chimpanzees in Kahuzi-Biega National Park, Zaire. In: McGrew WC, Marchant L, Nishida T, editors. *Great ape societies*. Cambridge: Cambridge University Press. p 82–98.
- Zihlman AL. 1997. Natural history of apes: life-history features of females and males. In: Morbeck ME, Galloay A, Zihlman A editors. *The evolving female: a life history perspective*. Princeton, NJ: Princeton University Press. p 185–197.
- Zihlman AL, McFarland RK. Submitted. Analysis of body mass in lowland gorillas. *Am J Phys Anthropol*.